

تم الرفع بواسطة  
م. معن ابو عيسى

# Calculus 1 Second

Palestine Technical University  
Mathematics Department.

Calculus I

Dr. Khalida Nazzal

Second Exam

Student #

Student name بالعربية:

معن سلام قعدان

Dec 7<sup>th</sup>, 2009

Sec #: RTN 11:00 - 12:00

## Part I: Choose the correct answer

(30 points)

1. The displacement of a particle moving in a straight line is given by  $s = \frac{t^4}{4} - 8t + 3$ .

When the velocity of the particle is equal to zero.

(a)  $t = 4$

(b)  $t = 0$

(c)  $t = 1$

(d)  $t = 2$

$s =$

$$\frac{t^3}{3} - 8$$

$$t^3 - 8 = 0$$

$$t^3 = 8 \Rightarrow t = \sqrt[3]{8} = 2$$

2. If the sides of a cube changes from 2 to 2.01, the new volume is approximately

(a) 0.12.

(b) 8.12

(c) 8.00

(d) 7.88.

$$dx = 0.01$$

$$8 + 0.12 = 8.12$$

$$V = x^3 = 2^3 = 8$$

$$dV = 3x^2 dx = 3(2)^2(0.01)$$

$$0.12$$

3. The equation of the tangent line to the curve  $x^3 - xy + y^3 = 7$  at  $P(2,1)$  is.

(a)  $5(y-1) = 11(x-2)$ .

(b)  $y = 11x - 21$ .

(c)  $(y-1) = -11(x-2)$ .

(d) else.

$$y-1 = \frac{13}{-5}(x-2)$$

$$y-1 = -\frac{13}{5}x + \frac{32}{5}$$

$$3x^2 + x \frac{dy}{dx} + y + 3y^2 \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = \frac{3(2)^2 + 1}{-2 - 3} = \frac{13}{-5}$$

$$3x^2 + y = -x \frac{dy}{dx} - 3y \frac{dy}{dx}$$

$$3x^2 + y = -(x + 3y) \frac{dy}{dx} \Rightarrow \frac{dy}{dx} = \frac{3x^2 + y}{-x - 3y}$$

4. Find the slope of the tangent line to the curve  $x = 2 + \sin t$ ,  $y = 3 + \cos t$ , at  $t = \frac{\pi}{4}$ .

(a) 1

(b) 0

(c) -1

(d) not defined

$$\frac{1}{\sqrt{2}} = x-2$$

$$\frac{1}{\sqrt{2}} + \frac{2}{1} = \frac{2\sqrt{2}+1}{\sqrt{2}}$$

$$\sin t = x-2$$

$$\cos t = y-3$$

$$\sin^2 t + \cos^2 t = 1$$

$$(x-2)^2 + (y-3)^2 = 1$$

$$(x-2)(x-2)$$

$$x^2 - 2x - 2x + 4$$

$$x^2 - 4x + 4$$

$$y^2 - 6y + 9$$

$$x^2 + 4x + 4 + y^2 - 6y + 9 = 1$$

5. The normal line at any point of the circle  $x^2 + y^2 = a^2$

(a) passes through the origin.

(b) has a negative slope

(c) is never a vertical line.

(d) is never a horizontal line.

$$2(x-2) + 3(y-3) = 0$$

$$2x - 4 + 3y - 9 = 0$$

$$2x - 4 = 3y - 9$$




$$y' (3 - 3y) = 2x - 4$$

$$y' = \frac{2x-4}{3-3y}$$

$$y' = 2\left(\frac{1}{\sqrt{2}} + 2\right) - 4$$

$$3 - 3\left(\frac{1}{\sqrt{2}} + 3\right)$$

$$\frac{\sqrt{2} + 4 - 4}{3 - \frac{3}{\sqrt{2}} - 9}$$





$$x = -1 \quad y = 2$$

6. Let  $x$  and  $y$  be functions of time  $t$  such that  $x^2 + y^3 = 9$ .

If  $\frac{dx}{dt} = 5$ , find  $\frac{dy}{dt}$  when  $x = -1$ .

- (a)  $\frac{5}{8}$   
 (b)  $\frac{5}{12}$   
 (c)  $\frac{1}{2\sqrt{2}}$   
 (d)  $\frac{5}{6}$

$$2x \frac{dx}{dt} + 3y^2 \frac{dy}{dt} = 0$$

$$2(-1)(5) + 3y^2 \frac{dy}{dt} = 0$$

$$-10 = 3y^2 \frac{dy}{dt}$$

$$x^2 = 9 - y^3$$

$$1 = 9 - y^3$$

$$y^3 = 9 - 1 = 8$$

$$y = 2$$

$$2(-1)(5) + 3(4) \frac{dy}{dt} = 0$$

$$-10 + 12 \frac{dy}{dt} = 0$$

$$10 = 12 \frac{dy}{dt} \Rightarrow \frac{dy}{dt} = \frac{10}{12} = \frac{5}{6}$$

7. Given that  $f(x) = 5 - \frac{4}{x}$ ,  $x \in [2, 4]$ . Find the value(s) of  $c$  in the conclusion of the Mean Value Theorem.

- (a)  $2\sqrt{2}, -2\sqrt{2}$   
 (b) 1  
 (c)  $2\sqrt{2}$   
 (d) 2, -2

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$5 - 4x^{-1}$$

$$= 4x^{-2} = \frac{4}{x^2}$$

$$= \frac{4}{x^2}$$

$$\frac{4}{c^2} = \frac{4 - 3}{4 - 2} \Rightarrow \frac{4}{c^2} = \frac{1}{2}$$

$$c^2 = 8$$

$$c = \pm\sqrt{8}$$

$$\pm 2\sqrt{2}$$

8. If  $h(x) = g(f(x))$  and  $f(2) = 5$ ,  $g(5) = 3$ ,  $f'(2) = 7$ , and  $g'(5) = -2$ , find  $h'(2)$ .

- (a) -14  
 (b) 21  
 (c) -10  
 (d) 14

$$h'(2) = g'(f(2)) = g'(5)$$

$$h'(2) = g'(f(2)) f'(2)$$

$$h'(2) = g'(5) (7) = -2 (7) = -14$$

9. A parametrization of the left half of the parabola  $y = x^2 + 2x$  could be


- (a)  $x = t^2$ ,  $y = t^4 + 2t^2$ ,  $-\infty < t < \infty$ .  
 (b)  $x = t$ ,  $y = t^2 + 2t$ ,  $-\infty < t < \infty$ .  
 (c)  $x = \sin t$ ,  $y = \sin^2 t + 2 \sin t$ ,  $-\infty < t < \infty$ .  
 (d)  $x = t$ ,  $y = t^2 + 2t$ ,  $t \leq -1$

$$x = t^2$$

$$y = (t^2)^2 + 2t^2$$

$$t^4 + 2t^2$$

$$(t^2)$$



10. The linearization of  $f(x) = \sqrt{x^2 + 8}$  at  $x = -1$  is

- (a)  $y = \frac{8}{3} - \frac{1}{3}x$   
 (b)  $y = 1 + \frac{1}{3}x$   
 (c)  $y = 1 + \frac{1}{2}x$   
 (d) None of the above

$$L(x) = f'(x_0) (x - x_0)$$

$$f'(x) = \frac{1}{2} (x^2 + 8)^{-\frac{1}{2}} (2x)$$

$$f'(x) = \frac{x}{\sqrt{x^2 + 8}}$$

$$= \frac{x}{\sqrt{x^2 + 8}}$$

$$3 + \frac{-1}{3} (x + 1)$$

$$3 - \frac{1}{3} (x + 1)$$

$$3 - \frac{1}{3} x - \frac{1}{3}$$

$$y =$$

$$2 - \frac{1}{3} x$$

$$y = 2 - \frac{1}{3}x$$



$$\frac{5}{2}x^{\frac{2}{3}} - x^{\frac{2}{3}}$$

$$x^{\frac{1}{3}} x^{\frac{2}{3}}$$

II. (20 points). Let  $f(x) = (\frac{5}{2} - x)x^{\frac{2}{3}}$ .

(a) find the intervals on which  $f$  is increasing or decreasing.

$$f'(x) = (\frac{5}{2} - x) \frac{2}{3} x^{-\frac{1}{3}} + x^{\frac{2}{3}} (-1) = \frac{10}{6} x^{-\frac{1}{3}} - \frac{2}{3} x^{\frac{2}{3}} - x^{\frac{2}{3}} = \frac{5}{3} x^{-\frac{1}{3}} - \frac{5}{3} x^{\frac{2}{3}}$$

$$= \frac{5}{3} \left( \frac{1}{\sqrt[3]{x}} - \sqrt[3]{x^2} \right) = \frac{5}{3} \left( \frac{1-x}{\sqrt[3]{x}} \right) = 0$$

$$\Rightarrow \frac{1-x}{\sqrt[3]{x}} = 0 \Rightarrow x=1, \text{ DNE } x=0$$

increasing  $0 < x < 1$   
decreasing  $x > 1, x < 0$   
 $(1, \infty) \cup (-\infty, 0)$

(b) find the x-coordinates of the local extrema of  $f$ .

$$f(0) = 0 \Rightarrow (0, 0) \Rightarrow \text{local minimum}$$

$$f(1) = \frac{3}{2}, (1, 1.5) \Rightarrow \text{local maximum}$$

(c) find the intervals on which  $f$  is CU or CD.

$$f'(x) = \frac{5}{3} \left( \frac{1-x}{\sqrt[3]{x}} \right) \Rightarrow f''(x) = \frac{\frac{3}{\sqrt[3]{x}} (-1) - (1-x) \left( \frac{1}{3} x^{-\frac{4}{3}} \right)}{x^{\frac{2}{3}}}$$


$$= \frac{-\frac{3}{\sqrt[3]{x}} - \left[ \frac{1}{3} x^{-\frac{1}{3}} - \frac{1}{3} x^{\frac{2}{3}} \right]}{x^{\frac{2}{3}}} = \frac{-\frac{3}{\sqrt[3]{x}} - \frac{1}{3} x^{-\frac{1}{3}} + \frac{1}{3} x^{\frac{2}{3}}}{x^{\frac{2}{3}}}$$

$$= \frac{-x^{\frac{1}{3}} - \frac{1}{3} x^{\frac{1}{3}} + \frac{1}{3} x^{\frac{2}{3}}}{x^{\frac{2}{3}}} = \frac{-x^{\frac{1}{3}} - \frac{1}{3} x^{\frac{1}{3}} + \frac{1}{3} x^{\frac{2}{3}}}{x^{\frac{2}{3}}} = -x^{-\frac{1}{3}} - \frac{1}{3} x^{-\frac{1}{3}} + \frac{1}{3} x^{\frac{2}{3}}$$



$$= -\frac{1}{\sqrt[3]{x}} - \frac{1}{3\sqrt[3]{x}} + \frac{1}{3} = 0 \Rightarrow \text{DNE at } x=0$$

(d) find the x-coordinates of the points of inflection of  $f$ .

$$f(0) = 0 \Rightarrow (0, 0)$$



\* \* find the cusp points if exists  
(e) Sketch the graph of  $f$ .





III. A particle moves along the curve  $y = x^{\frac{3}{2}}$  in the first quadrant in such a way that its distance from the origin increases at the rate 11 units per second. Find  $\frac{dx}{dt}$  when  $x = 3$ . (10 points).

$$y = x^{\frac{3}{2}} \quad , \quad \frac{ds}{dt} = 11$$

$$\text{distance} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$\therefore \text{distance} = \sqrt{x^2 + y^2} = \sqrt{x^2 + (x^{\frac{3}{2}})^2}$$

$$\textcircled{*} \text{ when } x = 3 \Rightarrow y = 3^{\frac{3}{2}} = \sqrt{3^3} = \sqrt{27}$$

~~$$ds = \sqrt{x^2 + y^2} \Rightarrow \frac{1}{2}(x^2 + y^2)^{-\frac{1}{2}} (2x dx + 2y dy)$$~~

$$\therefore S = \sqrt{x^2 + 27} = (x^2 + 27)^{\frac{1}{2}}$$

$$\therefore \frac{ds}{dt} = \frac{1}{2} (x^2 + 27)^{-\frac{1}{2}} (2x dx)$$

$$\therefore 11 = \frac{1}{2} (3^2 + 27)^{-\frac{1}{2}} (2(3) dx)$$

$$11 = \frac{1}{2\sqrt{36}} \cdot (6 dx)$$

$$11 = \frac{1}{2\sqrt{36}} \cdot 6(dx)$$

$$11 = \frac{1}{2} (dx) \Rightarrow dx = \boxed{22 \text{ units/second}}$$